

1 **OsLRR-RLK1, an early responsive leucine-rich repeat receptor-like kinase,**
2 **initiates rice defense responses against a chewing herbivore**

3

4 Lingfei Hu^{1,2*}, Meng Ye^{1,2*}, Peng Kuai¹, Miaofen Ye¹, Matthias Erb² and Yonggen
5 Lou¹

6

7 ¹State Key Laboratory of Rice Biology & Ministry of Agriculture Key Lab of
8 Molecular Biology of Crop Pathogens and Insects, Institute of Insect Sciences,
9 Zhejiang University, Hangzhou 310058, China; ²Institute of Plant Sciences, University
10 of Bern, Bern 3013, Switzerland.

11

12 Author for correspondence:

13 *Yonggen Lou*

14 *Tel: +86 (0)571 88982622*

15 *Email: yglou@zju.edu.cn*

16

17 * These authors contributed equally to this work

18

19 **Heading:** OsLRR-RLK1 initiates rice defenses against herbivores

20

Total word count (excluding summary, references and legends):	6104	No. of figures:	7 (Figs 1, 7 in color)
Summary:	202	No. of Tables:	0
Introduction:	965	No. of Supporting Information files:	12 (Figs S1-S11; Table S1)
Materials and Methods:	2519		
Results:	1594		
Discussion:	1796		
Acknowledgements:	129		

21

Summary

- Plants are constantly exposed to a variety of environmental stresses, including herbivory. How plants perceive herbivores on a molecular level is poorly understood. Leucine-rich repeat receptor-like kinases (LRR-RLKs), the largest subfamily of RLKs, are essential for plants to detect external stress signals and may therefore also be involved in herbivore perception.
- Here, we employed RNA interference silencing, phytohormone profiling and complementation as well as herbivore resistance assays to investigate the requirement of an LRR-RLK for initiating rice (*Oryza sativa*)-induced defense against the chewing herbivore striped stem borer (SSB) *Chilo suppressalis*.
- We discovered a plasma membrane-localized LRR-RLK, OsLRR-RLK1, whose transcription is strongly up-regulated by SSB attack and treatment with oral secretions of *Spodoptera frugiperda*. *OsLRR-RLK1* acts upstream of mitogen-activated protein kinase (MPK) cascades, and positively regulates defense-related MPKs, and WRKY transcription factors. Moreover, *OsLRR-RLK1* is a positive regulator of SSB-, but not wound-elicited levels of jasmonic acid and ethylene, trypsin protease inhibitor activity and plant resistance towards SSB.
- OsLRR-RLK1 therefore plays an important role in herbivory-induced defenses of rice. Given the well documented role of LRR-RLKs in the perception of stress-related molecules, we speculate that OsLRR-RLK1 may be involved in the perception of herbivory-associated molecular patterns.

Key words: *Chilo suppressalis*; defense responses; ethylene; herbivory perception; jasmonic acid; leucine-rich repeat receptor-like kinase; plant-herbivore interactions; rice

Introduction

In response to herbivore attack, plants activate a wide array of defenses which can reduce herbivore damage, including the initiation of phosphorylation-dependent signaling cascades such as mitogen-activated protein kinase (MPK) cascades, induction of defense-related signaling molecule biosynthesis such as jasmonic acid

(JA), salicylic acid (SA) and ethylene (ET), and the accumulation of defensive compounds (Wu & Baldwin, 2010; Erb *et al.*, 2012; Schuman & Baldwin, 2016). In order to activate their defenses, plants can recognize herbivores through damage-associated molecular patterns (DAMPs) as general wounding cues and herbivore-associated molecular patterns (HAMPs) as herbivore-specific cues (Bonaventure, 2012; Acevedo *et al.*, 2015; Schmelz, 2015). Although the specific pattern recognition by plants is well understood for pathogens (Zipfel, 2014; Couto & Zipfel, 2016), and DAMP perception is being unravelled (Choi *et al.*, 2014; Tanaka *et al.*, 2014; Tripathi *et al.*, 2018), the molecular basis of HAMP perception remains largely unknown (Mithofer & Boland, 2008; Gilardoni *et al.*, 2011; Mescher & De Moraes, 2015; Schmelz, 2015; Schuman & Baldwin, 2016).

Leucine-rich repeat receptor-like kinases (LRR-RLKs) have been shown to play a fundamental role in pattern recognition and initiation of downstream responses (Meng & Zhang, 2013; Macho & Zipfel, 2014). LRR-RLKs are characterized by tandem repeats of LRR motifs in their extracellular domains as well as an intracellular serine/threonine kinase domain (Tor *et al.*, 2009). LRR-RLKs have been shown to be involved in plant responses to wounding (Brutus *et al.*, 2010), gamma irradiation (Park *et al.*, 2014), drought (Osakabe *et al.*, 2005), salt (de Lorenzo *et al.*, 2009), heat (Jung *et al.*, 2015) and pathogens (Song *et al.*, 1995; Gomez-Gomez & Boller, 2000). The flagellin-sensitive 2 (FLS2), for instance, can recognize a conserved 22 amino acid epitope (flg22) from bacterial flagellin by its 28 extracellular LRRs (Gomez-Gomez & Boller, 2000; Gomez-Gomez *et al.*, 2001). Similarly, the elongation factor Tu receptor (EFR) can bind to *N*-acetylated 18 amino acid epitope (elf18) of the bacterial elongation factor Tu (ET-Tu) (Kunze *et al.*, 2004). Xa21 in rice (*Oryza sativa*) confers resistance to *Xanthomonas oryzae* pv. *oryzae* via the recognition of the tyrosine-sulfated protein RaxX (Pruitt *et al.*, 2015). LRR-RLKs have also been associated with plant responses to herbivory. Arabidopsis *pepr1* (Pep receptor 1) *pepr2* double mutants for instance show a reduced accumulation of oral secretion (OS)-elicited JA, and a decreased resistance to *Spodopora littoralis* larvae (Klauser *et al.*, 2015). Moreover, AtBAK1 (brassinosteroid insensitive1-associated

receptor kinase 1) is required for green peach aphid (*Myzus persicae*) elicitor-mediated callose deposition and reactive oxygen species (ROS) induction (Prince *et al.*, 2014). Accordingly, *atabak1* mutants are less resistant to the pea aphid (*Acyrtosiphon pisum*) (Prince *et al.*, 2014). In addition, silencing *NaBAK1* in wild tobacco (*Nicotiana attenuata*) attenuates wound- and OS-elicited JA and JA-isoleucine (JA-Ile) levels, but does not affect MPK activity and herbivore performance (Yang *et al.*, 2011). Despite these findings implicating LRR-RLKs in plant responses to herbivory, the underlying molecular mechanisms remain largely unexplored. Furthermore, the role of LRR-RLKs in plant-herbivore interactions in monocotyledons, as well as their potential to increase resistance against chewing herbivores, is unknown.

MPK cascades link cell surface receptors, such as LRR-RLKs, with downstream signaling components (Rodriguez *et al.*, 2010; Meng & Zhang, 2013). Generally, the stimulated receptors initiate the MPK cascades. Once started, the active MPK kinase kinases (MPKKKs or MEKKs) can activate downstream MPK kinases (MPKKs or MEKs), which subsequently activate MPKs through phosphorylation (Pitzschke, 2015). Activated MPKs phosphorylate their substrates, most of which are transcription factors and enzymes, thereby triggering downstream responses (Pitzschke, 2015). In Arabidopsis, the YODA-MKK4/MKK5-MPK3/MPK6 cascade functions at downstream of ERECTA receptor in regulating plant growth and development (Meng *et al.*, 2012). The MEKK1-MKK1/MKK2-MPK4 and MEKK1-MKK4/MKK5-MPK3/MPK6 can regulate immune responses which are activated by FLS2 after perception of flg22 (Asai *et al.*, 2002; Kong *et al.*, 2012). However, whether MEKK1 acts upstream of MKK4/MKK5 remains controversial (Meng & Zhang, 2013). Moreover, in *N. attenuata*, *Manduca sexta* OS can activate NaMEK2 (ortholog of AtMKK4/AtMKK5), wound-induced protein kinase (WIPK) and SA-induced protein kinase (SIPK, orthologs of AtMPK3 and AtMPK6), which have been reported to be involved in herbivore-induced defense responses via JA signaling (Wu *et al.*, 2007; Hettenhausen *et al.*, 2015). Similarly, the rice

OsMEK4-OsMPK3/OsMPK6 cascade positively regulates the JA signaling pathway and resistance to rice herbivores (Wang *et al.*, 2013; Li *et al.*, 2015).

WRKY transcription factors act as activators or repressors in plant defensive signaling webs downstream of MPK cascades (Ishihama & Yoshioka, 2012). WRKYs can be regulated by MPKs at transcriptional and/or post-translational levels (Ishihama & Yoshioka, 2012; Chi *et al.*, 2013; Li *et al.*, 2015). OsWRKY53, for instance, is activated by OsMPK3 and OsMPK6 through transcriptional induction and phosphorylation, thereby conferring rice resistance to both pathogens and herbivores (Chujo *et al.*, 2014; Hu *et al.*, 2015).

Rice, the most widely consumed food crop, suffers heavily from insect pests (Chen *et al.*, 2011). The striped stem borer (SSB) *Chilo suppressalis*, for instance, can bore into and feed on rice stems and causes large annual yield losses (Chen *et al.*, 2011). SSB attack induces a wide variety of defensive signaling pathways including MPKs, WRKYs, JA, SA and ET, which, in turn regulate rice defense responses (Zhou *et al.*, 2009; Zhou *et al.*, 2011; Lu *et al.*, 2014; Hu *et al.*, 2015).

Here, we isolated an SSB-induced LRR-RLK gene, *OsLRR-RLK1*, and characterized the involvement of this gene in herbivore-induced defense responses in rice. *OsLRR-RLK1* encodes a plasma membrane-localized protein and responses differentially to external stimuli. Using a reverse genetics approach, we obtained rice lines (*ir-ldrr*) with reduced expression of this gene and showed that it can positively regulate defense-related MPKs, WRKYs as well as the levels of herbivore-induced JA and ET, which subsequently mediated the activity of defensive trypsin protease inhibitors (TrypPIs) and resistance to SSB. Our study reveals that *OsLRR-RLK1* is an early responsive component of herbivore-related signaling pathways.

Materials and Methods

Plants and insects

The rice (*Oryza sativa*) genotypes used in this study were cultivar Xiushui 110 wild-type (WT) and transgenic lines of *ir-ldrr* (in this study), *as-mpk3* (Wang *et al.*,

2013), *as-mpk6* (Li *et al.*, 2015), *as-aos1* (Hu *et al.*, 2015), *as-lox* (Zhou *et al.*, 2009),
as-acs2 (Lu *et al.*, 2014) and *as-ics1* (Wang, 2012). These genotypes were cultivated
hydroponically as described in Hu *et al.* (2015) with the following composition: 1.43
mM NH₄NO₃, 1 mM CaCl₂, 0.32 mM NaH₂PO₄·2H₂O, 0.51 mM K₂SO₄, 1.64 mM
MgSO₄·7H₂O, 7.58 μM MnCl₂·4H₂O, 15.11 μM H₃BO₃, 0.12 μM CuSO₄·5H₂O, 0.06
μM (NH₄)₆Mo₇O₂₄·4H₂O, 0.12 μM ZnSO₄·7H₂O, 28.49 μM FeCl₃·6H₂O and 56.63
μM citric acid monohydrate (C₆H₈O₇·H₂O). The pH of the nutrient solution was
adjusted to 4.5-5.0 (Yoshida *et al.*, 1976). Forty day-old seedlings were individually
transferred to 500 ml hydroponic plastic pots, and then used for experiments 3 to 4 d
after transplanting. Larvae of the SSB *Chilo suppressalis* larvae were originally
obtained from rice fields in Hangzhou, China, and reared as described by Hu *et al.*
(2015). All experiments of this study were repeated at least twice.

Isolation and characterization of *OsLRR-RLK1*

The full-length cDNA of *OsLRR-RLK1* was amplified by PCR. The primers LRR-F
(5'-TGCAGCAGGCGAGTTTCATGA-3') and LRR-R
(5'-CACAAAAAAGAGGGAACTAA-3') were designed based on the sequence of
OsLRR-RLK1 (accession no. Os06g47650). The PCR products were cloned into the
pEASY-blunt cloning vector (TransGen) and sequenced.

***OsLRR-RLK1* sequence analysis**

Structural domain prediction was performed with SMART (Simple Modular
Architecture Research Tool, <http://smart.embl-heidelberg.de>; (Schultz *et al.*, 1998;
Letunic *et al.*, 2015) and Pfam (<http://pfam.sanger.ac.uk>) databases. Prediction of
transmembrane domains was performed with TMHMM
(<http://www.cbs.dtu.dk/services/TMHMM/>) web servers. Prediction of signal peptides
was performed using SignalP 4.0 (<http://www.cbs.dtu.dk/services/SignalP>). Protein
mass was estimated by ExPASy (http://web.expasy.org/compute_pi/, default setting).

Subcellular localization assay

For subcellular localization, the open reading frame of *OsLRR-RLK1* without the termination code was inserted into the pH7YWG2 plasmid to produce the fused OsLRR-RLK1-enhanced yellow fluorescent protein (EYFP) protein (Karimi *et al.*, 2005). The constructed plasmid was transformed into *Agrobacterium tumefaciens* C5851, and co-infiltrated into *Nicotiana benthamiana* leaves with the C5851 containing mCherry plasma membrane marker plasmid (Nelson *et al.*, 2007) at optical density at 600 nm of 0.7: 0.7. Small living pieces of *N. benthamiana* leaves were assayed for fluorescence 72 h after agroinfiltration. EYFP and mCherry fluorescence were observed and photographed by confocal microscopy (Leica TCS SP5). Spot detection and quantification on confocal micrographs were determined by the ImageJ software with Plot Profile function (<https://imagej.nih.gov/ij/index.html>).

Plant Treatments

For SSB treatment, one pre-starved third-instar SSB larva was placed on the stem of each plant. Typically, SSB larva crawls toward to the bottom portion of stem and chews a hole to feed on the inner tissues of the plant (Fig. S1). The moment the larva started to chew a hole was defined as time point zero for time course experiments. To measure SSB-induced plant responses, 2 cm portions of the stems around the entry hole were harvested at different time points after infestation. Control plants were not infested, and the same stem portions were harvested for analysis (Zhou *et al.*, 2009). For mechanical wounding, the lower portion of plant stems (approximately 2 cm long) was individually pierced 200 times with a sterilized needle. This piercing treatment aimed at mimicking the tissue damage inflicted by SSB. The damaged sections were harvested in a similar manner as for SSB experiments. Control plants were not pierced, and the same stem portions were harvested (Zhou *et al.*, 2009). For OS treatments, we could not rely on SSB OS, as the larvae do not regurgitate. We therefore used *Spodoptera frugiperda* OS. *Spodoptera frugiperda* attacks rice in the field (Pantoja *et al.*, 1986; Stout *et al.*, 2009) and produces OS that contains well-known defense elicitors such as fatty acid conjugates (FACs) (Yoshinaga *et al.*, 2010; Bonaventure *et al.*, 2011). Plants were wounded as described, and 10 μ l of *S. frugiperda* OS was

immediately applied to the wound sites (W + *S. frugiperda* OS). OS was collected from third instar *S. frugiperda* larvae that had been feeding on rice leaves for 48 h, and diluted 1:1 in autoclaved Milli-Q water before use. Ten microliters Milli-Q water were applied to the wounds of control plants (W + water). For JA or SA treatments, plants were individually sprayed with 2 ml of JA (100 µg ml⁻¹) or SA solution (70 µg ml⁻¹) in 50 mM sodium phosphate buffer. Controls (Buf) were sprayed with 2 ml of the buffer (Zhou *et al.*, 2009). For MeJA complementation, plant stems were individually treated with 100 µg of MeJA in 20 µl of lanolin paste. For lanolin treatment (+lanolin), plants were treated similarly with 20 µl of pure lanolin (Hu *et al.*, 2015).

QRT-PCR

For QRT-PCR analysis, five independent biological samples were used. Total RNA was isolated using the SV Total RNA Isolation System (Promega, catalog no. Z3100). One microgram of each total RNA sample was reverse transcribed with the PrimeScript RT-PCR Kit (TaKaRa, catalog no. RR014A). The QRT-PCR assay was performed on CFX96 Real-Time system (Bio-Rad). Gene expression levels were calculated using a standard curve method (Wong & Medrano, 2005). Briefly, a linear standard curve was constructed using serial dilutions of a specific cDNA standard, and drawn by plotting the threshold cycle (Ct) against the log₁₀ of the serial dilutions. The relative transcript levels of the target genes in all unknown samples were then determined according to the standard curve. The rice actin gene *OsACTIN* (accession no. Os03g50885) was used as an internal standard to normalize the cDNA concentrations. Primer specificity was confirmed by agarose gel electrophoresis, melting curve analysis, and sequence verification of cloned PCR amplicons. Primer pair efficiency was determined using the above standard curve method and was found to be between 95% and 105%. The primers, amplification efficiency, TaqMan probe sequences used for TaqMan QRT-PCR (*Premix Ex Taq*[™] [Probe qPCR]; Takara, catalog no. RR390A), and primer sequences for SYBR Green-based QRT-PCR (*SYBR*[®]*Premix Ex Taq*[™] II [Tli RNaseH Plus]; Takara, catalog no. RR820A) are

shown in Table S1.

Phylogenetic Analysis

The program MEGA 6.0 was used for the phylogenetic analysis (Tamura *et al.*, 2013). The protein sequences were aligned using the ClustalW method in MEGA 6.0 (pairwise alignment: gap opening penalty 10, gap extension penalty 0.1; multiple alignment: gap opening penalty 10, gap extension penalty 0.2, protein weight matrix using Gonnet). The residue-specific and hydrophilic penalties were on, and the end gap separation and the use negative separation matrix were off. Gap separation distance was 4, and the delay divergence cutoff (percentage) was at 30. This alignment was then used to generate an unrooted tree with statistical tests (parameters for phylogeny reconstruction were neighbor-joining method [Saitou & Nei, 1987] and bootstrap [Felsenstein, 1985], $n = 1,000$, amino acid, Poisson model, rate among sites: uniform rates gaps/missing, data treatment: complete deletion, traditional tree without modification for graphics) with MEGA 6.0.

Generation and characterization of transgenic plants

A 298-bp cDNA fragment of *OsLRR-RLK1* was inserted into the pCAMBIA-1301 transformation vector to yield an RNA interference (RNAi) construct (Fig. S2). The vector was inserted into Xiushui 110 using *A. tumefaciens*-mediated transformation. The rice transformation, screening of homozygous T₂ plants and identification of the number of insertions followed the same method as described in Zhou *et al.* (2009). Two T₂ homozygous lines (ir-1 and ir-3) were used in subsequent experiments.

MPK activation detection

One-month-old plants of different genotypes were randomly assigned to SSB or wounding treatments (see earlier). Plant stems were harvested at 0, 15, and 30 min after treatments. Total proteins were extracted from pooled stems of five replicates at each time point using the method described by Wu *et al.* (2007). Forty micrograms of total proteins were separated by SDS-PAGE and transferred onto Bio Trace pure

nitrocellulose blotting membrane (PALL). Immunoblotting was performed using the method described previously (Hu *et al.*, 2015). The primary antibodies used were the plant-actin rabbit polyclonal antibody (EarthOx, catalog no. E021080), which is used as a loading control or the rabbit monoclonal anti-phospho-ERK1/2 (anti-pT-E-pY) antibody (Cell Signaling Technologies, catalog no. 4370), which is specific for the activated (phosphorylated) form of the p44/42 MPKs, when catalytically activated by phosphorylation at the Thr-x-Tyr (TXY) motif (Segui-Simarro *et al.*, 2005; Anderson *et al.*, 2011). As a loading control, actin was detected on a replicate blot. Chemiluminescence-based detection (Thermo Scientific, catalog no. 32109) was performed using horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (Thermo Scientific, catalog no. 31460). The signal intensities of MPKs and loading actin in the immunoblots were quantified by the ImageJ software as described (Wu & Jackson, 2018). The signal intensity of OsMPK3, OsMPK6 or loading actin for the WT sample at 0 min was set to 1. The relative activation or quantity of all other samples at each time point was expressed relative to the WT sample at 0 min.

JA, JA-Ile, SA, and ET analysis

Plants of different genotypes were randomly assigned to SSB or wounding treatments (see above). Plant stems were harvested at 0, 1.5 and 3 h after the start of the treatments. JA, JA-Ile and SA were extracted with ethyl acetate spiked with labeled internal standards ($^{13}\text{C}_2$ -JA, $^{13}\text{C}_6$ -JA-Ile and D-SA, each with 100 ng) and analyzed with HPLC-MS/MS system following the method as described in (Lu *et al.*, 2015). For ET analysis, infested and control plants were covered with sealed glass cylinders (diameter, 4 cm; height, 50 cm). ET levels were determined using the method described by (Lu *et al.*, 2006). Each treatment at each time interval was replicated five times.

Analysis of TrypPI activity

The stems of WT plants and transgenic lines were harvested with SSB treatment for 3 d. The TrypPI activity was measured using a radial diffusion assay as described by

(Jongsma *et al.*, 1994; van Dam *et al.*, 2001). Each treatment was replicated five times.

Herbivore resistance experiments

For SSB performance, freshly hatched SSB neonates were introduced to feed on different rice genotypes. Larval mass was measured 12 d after the start of the experiment. Thirty replicate plants from each line and treatment were used. To determine differences in the tolerance of plants to SSB attack, the different genotypes were individually infested with one third-instar SSB larva. The damage levels of plants were checked and photographs were taken.

Data analysis

Differences in transcript levels of genes, concentrations of JA, JA-Ile, SA, and ET, and herbivore performance in different treatments, lines, or treatment times were determined by analysis of variance (ANOVA). When needed, pairwise or multiple comparisons of Least Squares Means (LSMeans) were corrected using the False Discovery Rate (FDR) method (Benjamini & Hochberg, 1995). All analyses were conducted using R 3.2.2 (R Foundation for Statistical Computing, Vienna, Austria).

Accession Numbers

Sequence data from this article can be found in the Rice Annotation Project under accession numbers *OsLRR-RLK1* (Os06g47650), *OsWRKY70* (Os05g39720), *OsWRKY53* (Os05g27730), *OsWRKY45* (Os05g25770), *OsWRKY24* (Os01g61080), *OsWRKY33* (Os03g33012), *OsWRKY30* (Os08g38990), *OsMEK4* (Os2g54600), *OsMPK3* (Os03g17700), *OsMPK6* (Os06g06090), *OsHI-LOX* (Os08g39840), *OsAOS1* (Os03g55800), *OsICS1* (Os09g19734), *OsACS2* (Os04g48850), and *OsACTIN* (Os03g50885).

Results

***OsLRR-RLK1* is an herbivory induced LRR-RLK**

Using microarrays, we identified a putative LRR-RLK that was up-regulated after SSB infestation (Zhou *et al.*, 2011), and obtained its full-length cDNA by reverse transcription PCR. The cDNA nucleotide sequence contains an open reading frame (ORF) of 3201 bp encoding a predicted protein of 1066 amino acids with an estimated molecular mass of 116 kDa (Fig. S3). Analysis of the deduced amino acid sequence predicted the presence of an N-terminal extracellular region including a signal peptide and multiple LRR domains, a single transmembrane domain, and a C-terminal cytoplasmic serine/threonine domain (Fig. S3). Based on these characteristics, the gene was named *OsLRR-RLK1* (for *O. sativa* leucine-rich repeat receptor-like kinase 1).

Comparative analysis of the RLK families in Arabidopsis and rice showed that *OsLRR-RLK1* belongs to the LRR-Xb subfamily, cluster 45, clade JA (Shiu *et al.*, 2004). Its closest characterized homologs in Arabidopsis were identified as plant peptide containing sulfated tyrosine 1 receptor (PSY1R), phytosulfokine receptor 2, (PSKR2), phytosulfokine receptor 1 (PSKR1), and receptor like protein 2 (RLP2) (Fig.S4 and S5). PSY1R and PSKR1 have been reported to modify the immunity of Arabidopsis to pathogens (Igarashi *et al.*, 2012; Mosher *et al.*, 2013; Shen & Diener, 2013), and we therefore hypothesized that *OsLRR-RLK1* may also be involved in rice defenses.

To determine the subcellular localization of *OsLRR-RLK1*, its coding region was fused to enhanced yellow fluorescent protein (EYFP) at the N-terminal end, and then expressed in *N. benthamiana* leaves under the control of *CaMV 35S* promoter (*35S::OsLRR-RLK1::EYFP*). As the membrane-localized marker AtPIP2A (Nelson *et al.*, 2007), a fluorescent signal was observed at the plasma membrane (Fig. 1). This suggests that *OsLRR-RLK1* may contribute to signal transduction as a component of a receptor system in the plasma membrane.

To investigate the regulation of *OsLRR-RLK1*, we examined its expression levels upon different elicitation treatments using quantitative real-time (QRT)-PCR. Compared with basal mRNA levels in non-manipulated stems (Con), *OsLRR-RLK1*

transcript levels were rapidly and strongly increased upon SSB attack (Fig. 2a). Mechanical wounding also increased *OsLRR-RLK1* mRNA levels, but the induction was weaker and slower compared to SSB attack (Fig. 2a, b). Adding *S. frugiperda* OS to the wounds strongly enhanced wound-induced expression of *OsLRR-RLK1* (Fig. 2c). JA treatment only marginally induced the *OsLRR-RLK1* transcript levels, and SA treatment did not (Fig. 2d, and Fig. S6). These data show that *OsLRR-RLK1* is strongly induced by herbivory, and responds strongly to OS and weakly to wounding alone.

***OsLRR-RLK1* silencing by RNA interference**

To study the function of *OsLRR-RLK1* in herbivore-induced responses in rice, transformed rice plants with reduced expression levels of *OsLRR-RLK1* were generated by *Agrobacterium tumefaciens*-based plant transformation. Two homozygous single insertion *OsLRR-RLK1*-silenced lines (*ir-lrr* lines: *ir-1* and *ir-3*) were selected and used to characterize the role of *OsLRR-RLK1* (Fig. S7). QRT-PCR analysis showed that both the constitutive and SSB-induced transcript levels of *OsLRR-RLK1* in *ir-lrr* lines were reduced 70-80% compared to wild-type (WT) plants (Fig. S8a). The RNAi construct did not co-silence the transcript levels of the genes whose nucleotide sequences have the highest similarity to *OsLRR-RLK1*, e. g. LOC_Os06g47760 (Top identity 92.72%, Top query coverage, 56.66%, rice genome annotation project algorithm), LOC_Os02g05960 (82.42%, 18.23%), LOC_Os02g05980 (82.37%, 18.89%), LOC_Os02g05920 (82.21%, 12.91%), and LOC_Os02g05940 (82.03%, 18.36%) (Fig. S8). The growth and morphology of *ir-lrr* lines were indistinguishable from those of WT plants at all the development stages both in the greenhouse and the field (Fig. S9).

***OsLRR-RLK1* regulates SSB-elicited OsMEK4, OsMPK3 and OsMPK6**

MPKs are required for rice defense in response to SSB attack (Wang *et al.*, 2013). To determine whether the silencing of *OsLRR-RLK1* changes MPK cascades, we measured the activation and expression levels of *OsMPK3* (also called *OsMPK5*) and

OsMPK6 (*OsMPK1* and *OsSIPK*) in WT and *ir-ldrr* plants. *OsMPK3* is the ortholog of *AtMPK3* and *WIPK*, and *OsMPK6* is the ortholog of *AtMPK6* and *SIPK* (Xie *et al.*, 2014). Their activation was determined by immunoblot analysis using an anti-phosphoERK1/2 (anti-pT-E-pY) antibody. This antibody specifically recognizes the phosphorylated residues within MPK activation loop (the so called pT-E-pY motif, where p denotes the phosphorylated residue), which is required for kinase activity. (Segui-Simarro *et al.*, 2005; Anderson *et al.*, 2011; Schwessinger *et al.*, 2015). In WT plants, SSB infestation rapidly and strongly induced the activation of *OsMPK3* and *OsMPK6*. The activation was reduced in *ir-ldrr* lines (Fig. 3a, and Fig. S10). Furthermore, SSB infestation rapidly and strongly induced the expression levels of *OsMPK3* and *OsMEK4*, and marginally induced the expression of *OsMPK6* in WT plants (Fig. 3b-d). The expression levels of *OsMPK3* and *OsMEK4* were significantly decreased in *ir-ldrr* plants compared with those in WT plants, whereas *OsMPK6* expression was not affected (Fig. 3b-d). To investigate whether *OsLRR-RLK1* is regulated by MPKs, *OsLRR-RLK1* expression was measured in *MPK*-silenced plants (*as-mpk3* and *as-mpk6*, Wang *et al.*, 2013; Li *et al.*, 2015). *OsLRR-RLK1* expression did not differ between WT and *MPK*-silenced plants (Fig. 3e, f). These results show that *OsLRR-RLK1* is a positive regulator of MPKs, and probably acts upstream of the MPK signaling pathway.

***OsLRR-RLK1* regulates defense-related WRKYs**

WRKYs are an important family of transcription factors to regulate plant defenses (Bakshi & Oelmüller, 2014). We have identified that *OsWRKY70*, *OsWRKY53*, *OsWRKY45*, *OsWRKY24* play crucial roles in the modulation of rice defense in response to herbivory (Li, 2012; Hu *et al.*, 2015; Li *et al.*, 2015; Huangfu *et al.*, 2016). Thus, we determined whether *OsLRR-RLK1* regulates the transcript levels of these four WRKYs and two additional defense-related WRKYs: *OsWRKY30* and *OsWRKY33* (Koo *et al.*, 2009; Han *et al.*, 2013). Silencing of *OsLRR-RLK1* greatly attenuated transcript accumulations of *OsWRKY70*, *OsWRKY53*, *OsWRKY45* and *OsWRKY24*, while it significantly enhanced *OsWRKY30* and *OsWRKY33* transcript

levels after infestation with SSB larvae for 15 and 30 min (Fig. 4).

OsLRR-RLK1 regulates SSB-elicited JA, SA and ET biosynthesis

Given that JA, JA-Ile, SA and ET are central mediators of rice defenses against herbivores (Zhou *et al.*, 2009; Zhou *et al.*, 2011; Lu *et al.*, 2014), we tested whether the reduced expression of *OsLRR-RLK1* alters the production of these phytohormones. JA and JA-Ile induction were significantly reduced in *ir-lrr* lines relative to WT plants (Fig. 5a, b). The transcript levels of JA biosynthesis genes *OsHI-LOX* and *OsAOS1* (Zhou *et al.*, 2009; Hu *et al.*, 2015) were also reduced in *ir-lrr* lines (Fig. 5c, d). ET also accumulated in smaller amounts in SSB-infested *ir-lrr* lines (Fig. 5e), which was associated with reduced expression of the ET biosynthetic gene *OsACS2* (Fig. 5f; Lu *et al.*, 2014). By contrast, *ir-lrr* lines accumulated significantly higher SSB-induced SA levels (Fig. 5g) and showed higher expression of the SA biosynthesis gene *OsICS1* (Wang, 2012) (Fig. 5h).

Most LRR-RLKs act upstream of hormonal signaling pathways (Antolin-Llovera *et al.*, 2012). To determine if this is also the case for OsLRR-RLK1 in rice, we quantified the expression of *OsLRR-RLK1* in transgenic plants with impaired JA, SA or ET biosynthesis (*as-lox*, Zhou *et al.*, 2009; *as-aos1*, Hu *et al.*, 2015; *as-ics1*, Wang, 2012; *as-acs2*, Lu *et al.*, 2014). The levels of constitutive and induced *OsLRR-RLK1* transcripts in *as-lox*, *as-aos1*, *as-ics1* and *as-acs2* lines were similar to those in WT plants (Fig. 6). Taken together, these results show that OsLRR-RLK1 acts upstream of JA, SA and ET signaling, and regulates the herbivory-induced biosynthesis of these hormones.

OsLRR-RLK1 does not regulate wound-elicited OsMPK3 and OsMPK6 activation and the levels of JA and SA

To further clarify the OsLRR-RLK1 regulation of herbivory-induced defense responses, we analyzed the MPK activation, JA and SA levels, in *ir-lrr* lines and WT plants after mechanical wounding. OsMPK3 was strongly activated at 30 min, while OsMPK6 was slightly induced at 15 min and decreased at 30 min by wounding.

However, in contrast with SSB infestation, the wound-induced MPK activation remained unchanged in *ir-lrr* lines relative to WT plants (Fig. S11a). Similarly, mechanical wounding significantly induced JA, JA-Ile and SA, but the induction of these phytohormones did not differ between *ir-lrr* lines and WT plants (Fig. S11b-d). These results suggest that OsLRR-RLK1 does not regulate wound-induced OsMPK3 and OsMPK6 activation and the JA, JA-Ile and SA production in the absence of an actual herbivore.

Silencing *OsLRR-RLK1* leads to decreased TrypPI activity and rice resistance to SSB

TrypPIs in rice are antidiigestive proteins which are strongly induced by SSB and slow down SSB growth (Zhou *et al.*, 2009). To analyze the function of *OsLRR-RLK1* in regulating TrypPIs, we determined the TrypPI activity in *ir-lrr* lines and WT plants 3d after SSB infestation. Compared with WT plants, *ir-lrr* lines showed a decrease of TrypPI activity of 45% (Fig. 7a). Consistently, SSB neonates gained more weight on *ir-lrr* lines than WT plants (Fig. 7b). Furthermore, *ir-lrr* lines were more susceptible to SSB than WT plants: after infestation by a third instar SSB larva for 7d, *ir-lrr* plants had completely died, whereas WT plants only showed mild dead heart symptoms (Fig. 7g).

To determine if the impaired TrypPI activity and rice resistance in *ir-lrr* lines can be rescued by restoring JA-dependent defenses, we treated *ir-lrr* plants with 100 µg methyl jasmonate (MeJA) in lanolin paste. This complementation restored the TrypPI activity to WT levels (Fig. 7c). Meanwhile, SSB larvae feeding on MeJA-treated *ir-lrr* plants exhibited the same performance as the ones feeding on WT plants (Fig. 7d). Moreover, in another experiment, we found that application of pure lanolin did not impair the difference in TrypPI activity and larval performance between *ir-lrr* lines and WT plants (Fig. 7 e and f). These results suggest that the compromised resistance of *OsLRR-RLK1*-silenced plants is a result of reduced JA signaling that leads to a reduction in defense activation, including TrypPI activity.

Discussion

This study identifies OsLRR-RLK1 as an early herbivore-responsive receptor-like kinase that is required for the initiation of rice defenses against a chewing herbivore.

RLKs can be classified on the basis of their extracellular domains (Tor *et al.*, 2009). OsLRR-RLK1 is placed into the LRR-RLK family due to its putative LRRs in the ectodomain. OsLRR-RLK1 shows high sequence similarity to three receptors PSY1R, PSKR2, PSKR1 and one LRR-receptor like protein (RLP) RLP2 in Arabidopsis (Fig. S5). PSY1R and PSKR1 can specifically bind their ligands, the tyrosine-sulfated peptides PSK or PSY1, via LRR domains (Matsubayashi *et al.*, 2002). PSKR2 is the paralog of PSKR1. RLP2 shares high sequence similarity with the receptor CLAVATA2 (CLV2), which can bind the small signaling peptide CLV3 (Wang *et al.*, 2010). Like PSKR1 and RLP2, OsLRR-RLK1 localized at the plasma membrane (Fig. 1). It is therefore plausible that OsLRR-RLK1-LRR binds to early signaling elements that are associated with herbivory, including for instance HAMPs themselves. Identifying the ligands of OsLRR-RLK1 is an exciting prospect of this work.

Plants can specifically distinguish HAMPs and DAMPs to tailor their defense responses (Bonaventure, 2012). In *N. attenuata*, *NaBAK1* transcript levels are quickly and strongly increased after *M. sexta* OS treatment, but only marginally increased by wounding (Yang *et al.*, 2011). In Arabidopsis, the application of OS as well as *S. littoralis* feeding strongly activates the promoters of *PEPR1* and *PEPR2*, whereas wounding alone does not (Klauser *et al.*, 2015). In our study, the transcript levels of *OsLRR-RLK1* were low in non-manipulated WT plants, but rapidly induced at the early stage (at 0.5 h) and strongly induced at the late stage (after 4 h) by SSB attack. The induction by larval OS was much stronger than mechanical wounding alone. Furthermore, *OsLRR-RLK1* regulated SSB-elicited, but not wounding-elicited MPK activation and phytohormone biosynthesis (Fig. 5, and Fig. S11). These results show that *OsLRR-RLK1* specifically responds to herbivory, and regulates herbivory-induced plant defenses.

Our work places the transcriptional induction of *OsLRR-RLK1* upstream of MPK,

WRKY and phytohormone signaling. Exogenous JA only marginally induced *OsLRR-RLK1* expression, and SA did not induce the expression of the gene at all (Fig. 2). Furthermore, impairing MPK, JA, SA, or ET signaling did not influence *OsLRR-RLK1* induction (Figs. 3 and 6). Thus, the rapid transcriptional induction of *OsLRR-RLK1* occurs independently of MPKs, JA, SA and ET. For instance it is possible that OsLRR-RLK1 activation triggers transcription via a positive feedback loop. In Arabidopsis, *PEPR1* and *PEPR2* are transcriptionally induced by small peptides (*AtPeps*), which are produced from damage-/herbivore-responsive *Precursor Protein of Plant Elicitor Peptide (PROPEP)* genes, which are in turn regulated by *AtWRKY33* (Huffaker *et al.*, 2006; Yamaguchi *et al.*, 2010; Logemann *et al.*, 2013). Furthermore, WRKY proteins can directly bind the W-box elements in the promoter of *RLK4* gene to regulate its expression (Du & Chen, 2000). Therefore, the transcriptional induction of *OsLRR-RLK1* by herbivory or wounding may be achieved through yet unidentified WRKY activity.

Plant MPK cascades play central roles in amplifying and transducing signals generated by receptors (Meng & Zhang, 2013). In Arabidopsis, for example, *pepr1pepr2* double mutants have markedly reduced expression levels of *MPK3* (Yamaguchi *et al.*, 2010). A loss of *SERK3/BAK1* results in a marked reduction of flg22 and elf18-triggered activation of MPK3 and MPK6 (Heese *et al.*, 2007), and the knock out mutants for *chitin elicitor receptor kinase 1 (CERK1)* completely lose the ability to activate MPK3 and MPK6 in response to chitin (Miya *et al.*, 2007). Respective CLV receptors possess unique activities for the regulation of MPK6 in Arabidopsis and *N. benthamiana* (Betsuyaku *et al.*, 2011). Here, we found that *OsMPK6* had high constitutive transcript levels and was only slightly induced by SSB infestation, while *OsMPK3* exhibited the opposite effect. Moreover, silencing *OsLRR-RLK1* reduced the expression levels of *OsMEK4* and *OsMPK3*, as well as the activation of OsMPK3 and OsMPK6 (Fig. 3). These data suggest that *OsMPK3* and *OsMPK6* might also be a pair of paralogous genes, like *AtMPK3* and *AtMPK6* in Arabidopsis (Menges *et al.*, 2008), and that OsLRR-RLK1 can activate MPK components upstream of OsMPK3 and OsMPK6. So far, several receptor-MPK

cascades have been reported. For example, cascades composed of CERK1-PBL27-MAPKKK5-MKK4/MKK5-MPK3/MPK6 in Arabidopsis and OsCERK1-OsRLCK185-OsMAPKKK18 (or OsMAPKKKε) -OsMKK4-OsMPK3/OsMPK6 in rice have recently been reported to be involved in chitin signaling (Yamada *et al.*, 2016; Wang *et al.*, 2017; Yamada *et al.*, 2017). Further researches should elucidate which MPK cascades function downstream of OsLRR-RLK1.

MPKs are known to be upstream regulators of WRKY transcription factors, including the ones that are regulated by OsLRR-RLK1 (Fig. 4). It has been reported that OsWRKY70, OsWRKY53, OsWRKY45, OsWRKY33, OsWRKY30 and OsWRKY24 are downstream of MPK cascades (Koo *et al.*, 2009; Li, 2012; Shen *et al.*, 2012; Ueno *et al.*, 2013; Chujo *et al.*, 2014; Hu *et al.*, 2015; Li *et al.*, 2015). OsWRKY70, OsWRKY53, OsWRKY45 and OsWRKY30 can physically interact with and be phosphorylated by OsMPK3 and/or OsMPK6 (Shen *et al.*, 2012; Ueno *et al.*, 2013; Chujo *et al.*, 2014; Hu *et al.*, 2015; Li *et al.*, 2015). In Arabidopsis, after perception by FLS2, flg22 induces WRKY22 and WRKY29 through activation of a MPK cascade composed of MEKK1, MKK4/MKK5, and MPK3/MPK6 (Asai *et al.*, 2002). In rice, upon herbivore or pathogen infestation, OsWRKY53 and OsWRKY70 are phosphorylated and activated by the OsMKK4-OsMPK3/OsMPK6 cascade (Chujo *et al.*, 2014; Li *et al.*, 2015). Therefore, the regulation of defense-related WRKYs probably occurs through MPK cascade which is modulated by OsLRR-RLK1. Additionally, the induction of some WRKYs as well as MPKs and hormone biosynthesis-related genes seems to be delayed after *OsLRR-RLK1*-silencing. This may be caused by functional redundancy with other homologous *RLK* genes or non-complete silencing of *OsLRR-RLK1*.

In *N. attenuata*, NaBAK1 regulates the accumulation of JA in responses to *M. sexta* (Yang *et al.*, 2011). In Arabidopsis, the lack of *PEPR1/PEPR2* receptors leads to reduced production of JA and JA-Ile after the application of *S. littoralis* OS (Klauser *et al.*, 2015). Furthermore, PSKR1 and PSY1R modify plant immunity to pathogens via JA- and SA- mediated signaling pathways (Mosher *et al.*, 2013). Here, we found

that OsLRR-RLK1 positively regulated the production of SSB-elicited JA, ET as well as the transcript levels of their biosynthesis-related genes, such as *OsHI-LOX*, *OsAOS1* and *OsACS2*, whereas it negatively influenced the accumulation of SA after SSB infestation, including the transcript levels of SA biosynthesis-related gene *OsICS1* (Fig. 5). Interestingly, it was also observed that expression levels of *OsHI-LOX*, *OsAOS1* and *OsACS2* were initially reduced in *ir-lrr* lines, and then back to WT levels by 90 min after SSB attack, while *OsICS1* showed the opposite effect. This may reflect the antagonistic crosstalk of JA/ET and SA signaling pathways in rice as reported previously (Lee *et al.*, 2004; Qiu *et al.*, 2007; Yuan *et al.*, 2007). Extensive studies have shown that MPKs and WRKYs mediate the biosynthesis of JA, SA, and ET in rice. For example, OsMPK3 positively regulates SSB-elicited JA levels (Wang *et al.*, 2013). OsMPK6 is involved in pathogen-related JA, SA accumulation (Shen *et al.*, 2010). OsWRKY70, OsWRKY53, OsWRKY45, and OsWRKY24 are implicated in herbivore-induced JA, SA and ET biosynthesis (Li, 2012; Hu *et al.*, 2015; Li *et al.*, 2015; Hu *et al.*, 2016; Huangfu *et al.*, 2016). OsWRKY33 and OsWRKY30 function as positive regulators of SA signaling pathway in rice (Koo *et al.*, 2009; Han *et al.*, 2013). Given the strong effects of OsLRR-RLK1 on MPKs and WRKYs found here, the regulation of JA, SA, and ET levels by OsLRR-RLK1 may be achieved mainly through MPK cascades and WRKYs.

In Arabidopsis, *pepr1pepr2* double mutants display reduced resistance to *S. litoralis* (Klauser *et al.*, 2015), and *bak1* mutant plants are compromised in immunity to aphids (Prince *et al.*, 2014). Here our experiments show that silencing of *OsLRR-RLK1* decreased the TrypPIs activity and the resistance of rice to SSB larvae, possibly via the impaired JA signaling (Fig. 7). This finding is consistent with our previous results showing that *as-lox* plants, which had lower elicited JA levels, were susceptible to SSB attack (Zhou *et al.*, 2009). Previous studies have also demonstrated that the ET signaling pathway positively regulates rice resistance to SSB: antisense expression of *OsACS2* (*as-acs2*) reduced herbivore-induced ET emission and the resistance of rice to SSB (Lu *et al.*, 2014). Therefore, we propose that the compromised resistance of *ir-lrr* lines to SSB is a result of low JA and ET levels,

which are positively mediated by OsLRR-RLK1.

In summary, our results demonstrate that OsLRR-RLK1 functions as a potential herbivore-recognition receptor of rice, and initiates induced defenses against SSB. We propose that the membrane-localized OsLRR-RLK1 may either directly bind to HAMPs or indirectly bind to other HAMP-induced early signaling molecules and immediately activate MPKs, which subsequently increase the activity of downstream WRKYs. Then, the activated MPKs and WRKYs regulate the biosynthesis of herbivore-related phytohormones, including JA, SA and ET, which result in effective induced defense responses against SSB. Our findings show how a plant employs an early responsive LRR-RLK to trigger specific defense responses against herbivores. We propose OsLRR-RLK1 as a candidate receptor of early signaling molecules that are associated with herbivory.

Acknowledgments

We thank Wei Huang and Maxime Hervé for their assistance with statistical analyses. The study was jointly sponsored by the key program of National Natural Science Foundation of China (31330065), the Fund for Agro-scientific Research in the Public Interest (201403030), the National Natural Science Foundation of China (31272052) and the earmarked fund for China Agriculture Research System (CARS-01-40). The work of L. H., M. Y., M. E. and Y. L. done in Switzerland was funded by the Swiss National Science Foundation (Grant 155781 to M. E.), the Sino-Swiss Science and Technology Cooperation (Exchange Grant Nr. EG 03-032016 to M. E. and Y. L.) and the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC-2016-STG 714239 to M. E.). No conflict of interest declared.

Author Contributions

L. H., M. Y., M. E. and Y. L. designed the research; L. H., M. Y., P. K., and M. Y. performed experiments; L. H., M. Y., P. K., M. Y., M. E. and Y. L. analyzed and interpreted data; L. H., M. Y., M. E., and Y. L. prepared and wrote the article. All

623 authors read and approved the manuscript.

References

- Acevedo FE, Rivera-Vega LJ, Chung SH, Ray S, Felton GW. 2015. Cues from chewing insects - the intersection of DAMPs, HAMPs, MAMPs and effectors. *Current Opinion in Plant Biology* **26**: 80-86.
- Anderson JC, Bartels S, Gonzalez Besteiro MA, Shahollari B, Ulm R, Peck SC. 2011. Arabidopsis MAP Kinase Phosphatase 1 (AtMKP1) negatively regulates MPK6-mediated PAMP responses and resistance against bacteria. *Plant Journal* **67**: 258-268.
- Antolin-Llovera M, Ried MK, Binder A, Parniske M. 2012. Receptor kinase signaling pathways in plant-microbe interactions. *Annual Review of Phytopathology* **50**: 451-473.
- Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, Gomez-Gomez L, Boller T, Ausubel FM, Sheen J. 2002. MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* **415**: 977-983.
- Bakshi M, Oelmuller R. 2014. WRKY transcription factors: Jack of many trades in plants. *Plant Signaling & Behavior* **9**: e27700.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **57**: 289-300.
- Betsuyaku S, Takahashi F, Kinoshita A, Miwa H, Shinozaki K, Fukuda H, Sawa S. 2011. Mitogen-activated protein kinase regulated by the CLAVATA receptors contributes to shoot apical meristem homeostasis. *Plant Cell Physiology* **52**: 14-29.
- Bonaventure G. 2012. Perception of insect feeding by plants. *Plant Biology* **14**: 872-880.
- Bonaventure G, VanDoorn A, Baldwin IT. 2011. Herbivore-associated elicitors: FAC signaling and metabolism. *Trends in Plant Science* **16**: 294-299.
- Brutus A, Sicilia F, Macone A, Cervone F, De Lorenzo G. 2010. A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a

654 receptor of oligogalacturonides. *Proceedings of the National Academy of*
655 *Sciences, USA* **107**: 9452-9457.

656 **Chen M, Shelton A, Ye GY. 2011.** Insect-resistant genetically modified rice in China:
657 from research to commercialization. *Annual Review of Entomology* **56**:
658 81-101.

659 **Chi Y, Yang Y, Zhou Y, Zhou J, Fan B, Yu JQ, Chen Z. 2013.** Protein-protein
660 interactions in the regulation of WRKY transcription factors. *Molecular Plant*
661 **6**: 287-300.

662 **Choi J, Tanaka K, Cao Y, Qi Y, Qiu J, Liang Y, Lee SY, Stacey G. 2014.**
663 Identification of a plant receptor for extracellular ATP. *Science* **343**: 290-294.

664 **Chujo T, Miyamoto K, Ogawa S, Masuda Y, Shimizu T, Kishi-Kaboshi M,**
665 **Takahashi A, Nishizawa Y, Minami E, Nojiri H, et al. 2014.** Overexpression
666 of phosphomimic mutated OsWRKY53 leads to enhanced blast resistance in
667 rice. *PLoS One* **9**: e98737.

668 **Couto D, Zipfel C. 2016.** Regulation of pattern recognition receptor signalling in
669 plants. *Nature Reviews Immunology* **16**: 537-552.

670 **de Lorenzo L, Merchan F, Laporte P, Thompson R, Clarke J, Sousa C, Crespi M.**
671 **2009.** A novel plant leucine-rich repeat receptor kinase regulates the response
672 of *Medicago truncatula* roots to salt stress. *Plant Cell* **21**: 668-680.

673 **Du L, Chen Z. 2000.** Identification of genes encoding receptor-like protein kinases as
674 possible targets of pathogen- and salicylic acid-induced WRKY DNA-binding
675 proteins in *Arabidopsis*. *Plant Journal* **24**: 837-847.

676 **Erb M, Meldau S, Howe GA. 2012.** Role of phytohormones in insect-specific plant
677 reactions. *Trends in Plant Science* **17**: 250-259.

678 **Felsenstein J. 1985.** Confidence limits on phylogenies: an approach using the
679 bootstrap. *Evolution* **39**: 783-791.

680 **Gilardoni PA, Hettenhausen C, Baldwin IT, Bonaventure G. 2011.** *Nicotiana*
681 *attenuata* *LECTIN RECEPTOR KINASE1* suppresses the insect-mediated
682 inhibition of induced defense responses during *Manduca sexta* herbivory.
683 *Plant Cell* **23**: 3512-3532.

- Gomez-Gomez L, Bauer Z, Boller T. 2001.** Both the extracellular leucine-rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signaling in *Arabidopsis*. *Plant Cell* **13**: 1155-1163.
- Gomez-Gomez L, Boller T. 2000.** FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. *Molecular Cell* **5**: 1003-1011.
- Han M, Ryu HS, Kim CY, Park DS, Ahn YK, Jeon JS. 2013.** *OsWRKY30* is a transcription activator that enhances rice resistance to the *Xanthomonas oryzae* pathovar *oryzae*. *Journal of Plant Biology* **56**: 258-265.
- Heese A, Hann DR, Gimenez-Ibanez S, Jones AM, He K, Li J, Schroeder JI, Peck SC, Rathjen JP. 2007.** The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proceedings of the National Academy of Sciences, USA* **104**: 12217-12222.
- Hettenhausen C, Schuman MC, Wu J. 2015.** MAPK signaling: a key element in plant defense response to insects. *Insect Science* **22**: 157-164.
- Hu L, Ye M, Li R, Lou Y. 2016.** *OsWRKY53*, a versatile switch in regulating herbivore-induced defense responses in rice. *Plant Signaling & Behavior* **11**: e1169357.
- Hu L, Ye M, Li R, Zhang T, Zhou G, Wang Q, Lu J, Lou Y. 2015.** The rice transcription factor *WRKY53* suppresses herbivore-induced defenses by acting as a negative feedback modulator of mitogen-activated protein kinase activity. *Plant Physiology* **169**: 2907-2921.
- Huangfu J, Li J, Li R, Ye M, Kuai P, Zhang T, Lou Y. 2016.** The transcription factor *OsWRKY45* negatively modulates the resistance of rice to the brown planthopper *Nilaparvata lugens*. *International Journal of Molecular Sciences* **17**: 697.
- Huffaker A, Pearce G, Ryan CA. 2006.** An endogenous peptide signal in *Arabidopsis* activates components of the innate immune response. *Proceedings of the National Academy of Sciences, USA* **103**: 10098-10103.
- Igarashi D, Tsuda K, Katagiri F. 2012.** The peptide growth factor, phytosulfokine,

attenuates pattern-triggered immunity. *Plant Journal* **71**: 194-204.

Ishihama N, Yoshioka H. 2012. Post-translational regulation of WRKY transcription factors in plant immunity. *Current Opinion in Plant Biology* **15**: 431-437.

Jongsma MA, Bakker PL, Visser B, Stiekema WJ. 1994. Trypsin-inhibitor activity in mature tobacco and tomato plants is mainly induced locally in response to insect attack, wounding and virus-infection. *Planta* **195**: 29-35.

Jung CG, Hwang SG, Park YC, Park HM, Kim DS, Park DH, Jang CS. 2015. Molecular characterization of the cold- and heat-induced *Arabidopsis PXL1* gene and its potential role in transduction pathways under temperature fluctuations. *Journal of Plant Physiology* **176**: 138-146.

Karimi M, De Meyer B, Hilson P. 2005. Modular cloning in plant cells. *Trends in Plant Science* **10**: 103-105.

Klauser D, Desurmont GA, Glauser G, Vallat A, Flury P, Boller T, Turlings TC, Bartels S. 2015. The *Arabidopsis* Pep-PEPR system is induced by herbivore feeding and contributes to JA-mediated plant defence against herbivory. *Journal of Experimental Botany* **66**: 5327-5336.

Kong Q, Qu N, Gao M, Zhang Z, Ding X, Yang F, Li Y, Dong OX, Chen S, Li X, et al. 2012. The MEKK1-MKK1/MKK2-MPK4 kinase cascade negatively regulates immunity mediated by a mitogen-activated protein kinase kinase kinase in *Arabidopsis*. *Plant Cell* **24**: 2225-2236.

Koo SC, Moon BC, Kim JK, Kim CY, Sung SJ, Kim MC, Cho MJ, Cheong YH. 2009. OsBWMK1 mediates SA-dependent defense responses by activating the transcription factor OsWRKY33. *Biochemical and Biophysical Research Communications* **387**: 365-370.

Kunze G, Zipfel C, Robatzek S, Niehaus K, Boller T, Felix G. 2004. The N terminus of bacterial elongation factor Tu elicits innate immunity in *Arabidopsis* plants. *Plant Cell* **16**: 3496-3507.

Lee A, Cho K, Jang S, Rakwal R, Iwahashi H, Agrawal GK, Shim J, Han O. 2004. Inverse correlation between jasmonic acid and salicylic acid during early wound response in rice. *Biochemical and Biophysical Research*

744 *Communications* **318**: 734-738.

745 **Letunic I, Doerks T, Bork P. 2015.** SMART: recent updates, new developments and
 746 status in 2015. *Nucleic Acids Research* **43**: D257-260.

747 **Li R. 2012.** *Function characterization of herbivore resistance-related genes*
 748 *OsWRKY24, OsWRKY70 and OsNPR1 in rice*. PhD thesis, Zhejiang
 749 University, Hangzhou, China.

750 **Li R, Zhang J, Li J, Zhou G, Wang Q, Bian W, Erb M, Lou Y. 2015.** Prioritizing
 751 plant defence over growth through WRKY regulation facilitates infestation by
 752 non-target herbivores. *Elife* **4**: e04805.

753 **Logemann E, Birkenbihl RP, Rawat V, Schneeberger K, Schmelzer E, Somssich**
 754 **IE. 2013.** Functional dissection of the *PROPEP2* and *PROPEP3* promoters
 755 reveals the importance of WRKY factors in mediating microbe-associated
 756 molecular pattern-induced expression. *New Phytologist* **198**: 1165-1177.

757 **Lu J, Li J, Ju H, Liu X, Erb M, Wang X, Lou Y. 2014.** Contrasting effects of
 758 ethylene biosynthesis on induced plant resistance against a chewing and a
 759 piercing-sucking herbivore in rice. *Molecular Plant* **7**: 1670-1682.

760 **Lu J, Robert CA, Riemann M, Cosme M, Mene-Saffrane L, Massana J, Stout MJ,**
 761 **Lou Y, Gershenzon J, Erb M. 2015.** Induced jasmonate signaling leads to
 762 contrasting effects on root damage and herbivore performance. *Plant*
 763 *Physiology* **167**: 1100-1116.

764 **Lu Y, Wang X, Lou Y, Cheng J. 2006.** Role of ethylene signaling in the production
 765 of rice volatiles induced by the rice brown planthopper *Nilaparvata lugens*.
 766 *Chinese Science Bulletin* **51**: 2457-2465.

767 **Macho AP, Zipfel C. 2014.** Plant PRRs and the activation of innate immune signaling.
 768 *Molecular Cell* **54**: 263-272.

769 **Matsubayashi Y, Ogawa M, Morita A, Sakagami Y. 2002.** An LRR receptor kinase
 770 involved in perception of a peptide plant hormone, phytosulfokine. *Science*
 771 **296**: 1470-1472.

772 **Meng X, Wang H, He Y, Liu Y, Walker JC, Torii KU, Zhang S. 2012.** A MAPK
 773 cascade downstream of ERECTA receptor-like protein kinase regulates

774 *Arabidopsis* inflorescence architecture by promoting localized cell
 775 proliferation. *Plant Cell* **24**: 4948-4960.

776 **Meng X, Zhang S. 2013.** MAPK cascades in plant disease resistance signaling.
 777 *Annual Review of Phytopathology* **51**: 245-266.

778 **Mescher MC, De Moraes CM. 2015.** Role of plant sensory perception in
 779 plant-animal interactions. *Journal of Experimental Botany* **66**: 425-433.

780 **Mithofer A, Boland W. 2008.** Recognition of herbivory-associated molecular
 781 patterns. *Plant Physiology* **146**: 825-831.

782 **Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, Shirasu K, Narusaka Y,**
 783 **Kawakami N, Kaku H, Shibuya N. 2007.** CERK1, a LysM receptor kinase,
 784 is essential for chitin elicitor signaling in *Arabidopsis*. *Proceedings of the*
 785 *National Academy of Sciences, USA* **104**: 19613-19618.

786 **Mosher S, Seybold H, Rodriguez P, Stahl M, Davies KA, Dayaratne S, Morillo**
 787 **SA, Wierzba M, Favery B, Keller H, et al. 2013.** The tyrosine-sulfated
 788 peptide receptors PSKR1 and PSY1R modify the immunity of *Arabidopsis* to
 789 biotrophic and necrotrophic pathogens in an antagonistic manner. *Plant*
 790 *Journal* **73**: 469-482.

791 **Nelson BK, Cai X, Nebenfuhr A. 2007.** A multicolored set of *in vivo* organelle
 792 markers for co-localization studies in *Arabidopsis* and other plants. *Plant*
 793 *Journal* **51**: 1126-1136.

794 **Osakabe Y, Maruyama K, Seki M, Satou M, Shinozaki K, Yamaguchi-Shinozaki**
 795 **K. 2005.** Leucine-rich repeat receptor-like kinase1 is a key membrane-bound
 796 regulator of abscisic acid early signaling in *Arabidopsis*. *Plant Cell* **17**:
 797 1105-1119.

798 **Pantoja A, Smith CM, Robinson JF. 1986.** Evaluation of rice germ plasm for
 799 resistance to the fall armyworm (Lepidoptera, Noctuidae). *Journal of*
 800 *Economic Entomology* **79**: 1319-1323.

801 **Park S, Moon JC, Park YC, Kim JH, Kim DS, Jang CS. 2014.** Molecular
 802 dissection of the response of a rice leucine-rich repeat receptor-like kinase
 803 (LRR-RLK) gene to abiotic stresses. *Journal of Plant Physiology* **171**:

1645-1653.

Pitzschke A. 2015. Modes of MAPK substrate recognition and control. *Trends in Plant Science* **20**: 49-55.

Prince DC, Drurey C, Zipfel C, Hogenhout SA. 2014. The leucine-rich repeat receptor-like kinase BRASSINOSTEROID INSENSITIVE1-ASSOCIATED KINASE1 and the cytochrome P450 PHYTOALEXIN DEFICIENT3 contribute to innate immunity to aphids in Arabidopsis. *Plant Physiology* **164**: 2207-2219.

Pruitt RN, Schwessinger B, Joe A, Thomas N, Liu F, Albert M, Robinson MR, Chan LJ, Luu DD, Chen H, et al. 2015. The rice immune receptor XA21 recognizes a tyrosine-sulfated protein from a Gram-negative bacterium. *Science Advances* **1**: e1500245.

Qiu D, Xiao J, Ding X, Xiong M, Cai M, Cao Y, Li X, Xu C, Wang S. 2007. OsWRKY13 mediates rice disease resistance by regulating defense-related genes in salicylate- and jasmonate-dependent signaling. *Molecular Plant-Microbe Interactions* **20**: 492-499.

Rodriguez MC, Petersen M, Mundy J. 2010. Mitogen-activated protein kinase signaling in plants. *Annual Review of Plant Biology* **61**: 621-649.

Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**: 406-425.

Schmelz EA. 2015. Impacts of insect oral secretions on defoliation-induced plant defense. *Current Opinion in Insect Science* **9**: 7-15.

Schultz J, Milpetz F, Bork P, Ponting CP. 1998. SMART, a simple modular architecture research tool: identification of signaling domains. *Proceedings of the National Academy of Sciences, USA* **95**: 5857-5864.

Schuman MC, Baldwin IT. 2016. The layers of plant responses to insect herbivores. *Annual Review of Entomology* **61**: 373-394.

Schwessinger B, Bahar O, Thomas N, Holton N, Nekrasov V, Ruan D, Canlas PE, Daudi A, Petzold CJ, Singan VR, et al. 2015. Transgenic expression of the

dicotyledonous pattern recognition receptor EFR in rice leads to
ligand-dependent activation of defense responses. *PLoS Pathogens* **11**:
e1004809.

Segui-Simarro J, Testillano P, Jouannic S, Henry Y, Risueno M. 2005.
Mitogen-activated protein kinases are developmentally regulated during
stress-induced microspore embryogenesis in *Brassica napus* L. *Histochemistry
and Cell Biology* **123**: 541-551.

Shen H, Liu C, Zhang Y, Meng X, Zhou X, Chu C, Wang X. 2012. OsWRKY30 is
activated by MAP kinases to confer drought tolerance in rice. *Plant Molecular
Biology* **80**: 241-253.

Shen X, Yuan B, Liu H, Li X, Xu C, Wang S. 2010. Opposite functions of a rice
mitogen-activated protein kinase during the process of resistance against
Xanthomonas oryzae. *Plant Journal* **64**: 86-99.

Shen Y, Diener AC. 2013. *Arabidopsis thaliana* RESISTANCE TO FUSARIUM
OXYSPORUM 2 implicates tyrosine-sulfated peptide signaling in
susceptibility and resistance to root infection. *PLoS Genetics* **9**: e1003525.

Shiu SH, Karlowski WM, Pan RS, Tzeng YH, Mayer KFX, Li WH. 2004.
Comparative analysis of the receptor-like kinase family in Arabidopsis and
rice. *Plant Cell* **16**: 1220-1234.

**Song WY, Wang GL, Chen LL, Kim HS, Pi LY, Holsten T, Gardner J, Wang B,
Zhai WX, Zhu LH, et al. 1995.** A receptor kinase-like protein encoded by the
rice disease resistance gene, *Xa21*. *Science* **270**: 1804-1806.

Stout MJ, Riggio MR, Yang Y. 2009. Direct Induced Resistance in *Oryza sativa* to
Spodoptera frugiperda. *Environmental Entomology* **38**: 1174-1181.

Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. 2013. MEGA6: molecular
evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*
30: 2725-2729.

Tanaka K, Choi JM, Cao YR, Stacey G. 2014. Extracellular ATP acts as a
damage-associated molecular pattern (DAMP) signal in plants. *Frontiers in
Plant Science* **5**: 446.

- Tor M, Lotze MT, Holton N. 2009.** Receptor-mediated signalling in plants: molecular patterns and programmes. *Journal of Experimental Botany* **60**: 3645-3654.
- Tripathi D, Zhang T, Koo AJ, Stacey G, Tanaka K. 2018.** Extracellular ATP acts on jasmonate signaling to reinforce plant defense. *Plant Physiology* **176**: 511-523.
- Ueno Y, Yoshida R, Kishi-Kaboshi M, Matsushita A, Jiang CJ, Goto S, Takahashi A, Hirochika H, Takatsuji H. 2013.** MAP kinases phosphorylate rice WRKY45. *Plant Signaling & Behavior* **8**: e24510.
- van Dam NM, Horn M, Mares M, Baldwin IT. 2001.** Ontogeny constrains systemic protease inhibitor response in *Nicotiana attenuata*. *Journal of Chemical Ecology* **27**: 547-568.
- Wang B. 2012.** *Functional analysis of herbivore resistance-related genes OsICS and OsHPL3 in rice*. PhD thesis, Zhejiang University, Hangzhou, China.
- Wang C, Wang G, Zhang C, Zhu PK, Dai HL, Yu N, He ZH, Xu L, Wang ET. 2017.** OsCERK1-mediated chitin perception and immune signaling requires receptor-like cytoplasmic kinase 185 to activate an MAPK cascade in rice. *Molecular Plant* **10**: 619-633.
- Wang G, Long Y, Thomma BP, de Wit PJ, Angenent GC, Fiers M. 2010.** Functional analyses of the CLAVATA2-like proteins and their domains that contribute to CLAVATA2 specificity. *Plant Physiology* **152**: 320-331.
- Wang Q, Li J, Hu L, Zhang T, Zhang G, Lou Y. 2013.** OsMPK3 positively regulates the JA signaling pathway and plant resistance to a chewing herbivore in rice. *Plant Cell Reports* **32**: 1075-1084.
- Wong ML, Medrano JF. 2005.** Real-time PCR for mRNA quantitation. *Biotechniques* **39**: 75-85.
- Wu J, Baldwin IT. 2010.** New insights into plant responses to the attack from insect herbivores. *Annual Review of Genetics* **44**: 1-24.
- Wu J, Hettenhausen C, Meldau S, Baldwin IT. 2007.** Herbivory rapidly activates MAPK signaling in attacked and unattacked leaf regions but not between leaves of *Nicotiana attenuata*. *Plant Cell* **19**: 1096-1122.

894 **Wu Q, Jackson D 2018.** Detection of MAPK3/6 phosphorylation during
895 hypersensitive response (HR)-associated programmed cell death in plants. In:
896 De Gara L, Locato V eds. *Plant Programmed Cell Death: Methods and*
897 *Protocols*. New York, NY: Springer New York, 153-161.

898 **Xie K, Chen J, Wang Q, Yang Y. 2014.** Direct phosphorylation and activation of a
899 mitogen-activated protein kinase by a calcium-dependent protein kinase in rice.
900 *Plant Cell* **26**: 3077-3089.

901 **Yamada K, Yamaguchi K, Shirakawa T, Nakagami H, Mine A, Ishikawa K,**
902 **Fujiwara M, Narusaka M, Narusaka Y, Ichimura K, et al. 2016.** The
903 *Arabidopsis* CERK1-associated kinase PBL27 connects chitin perception to
904 MAPK activation. *EMBO Journal* **35**: 2468-2483.

905 **Yamada K, Yamaguchi K, Yoshimura S, Terauchi A, Kawasaki T. 2017.**
906 Conservation of chitin-induced MAPK signaling pathways in rice and
907 *Arabidopsis*. *Plant and Cell Physiology* **58**: 993-1002.

908 **Yamaguchi Y, Huffaker A, Bryan AC, Tax FE, Ryan CA. 2010.** PEPR2 is a second
909 receptor for the Pep1 and Pep2 peptides and contributes to defense responses
910 in *Arabidopsis*. *Plant Cell* **22**: 508-522.

911 **Yang DH, Hettenhausen C, Baldwin IT, Wu J. 2011.** BAK1 regulates the
912 accumulation of jasmonic acid and the levels of trypsin proteinase inhibitors in
913 *Nicotiana attenuata*'s responses to herbivory. *Journal of Experimental Botany*
914 **62**: 641-652.

915 **Yoshida S, Forno DA, Cock JH, Gomez KA. 1976.** *Laboratory manual for*
916 *physiological studies of rice*. Manila, Philippines: The International Rice
917 Research Institute.

918 **Yoshinaga N, Alborn HT, Nakanishi T, Suckling DM, Nishida R, Tumlinson JH,**
919 **Mori N. 2010.** Fatty acid-amino acid conjugates diversification in
920 lepidopteran caterpillars. *Journal of Chemical Ecology* **36**: 319-325.

921 **Yuan YX, Zhong SH, Li Q, Zhu ZR, Lou YG, Wang LY, Wang JJ, Wang MY, Li**
922 **QL, Yang DL, et al. 2007.** Functional analysis of rice *NPRI*-like genes
923 reveals that *OsNPRI/NH1* is the rice orthologue conferring disease resistance

924 with enhanced herbivore susceptibility. *Plant Biotechnology Journal* **5**:
925 313-324.

926 **Zhou G, Qi J, Ren N, Cheng J, Erb M, Mao B, Lou Y. 2009.** Silencing *OsHI-LOX*
927 makes rice more susceptible to chewing herbivores, but enhances resistance to
928 a phloem feeder. *Plant Journal* **60**: 638-648.

929 **Zhou G, Wang X, Yan F, Wang X, Li R, Cheng J, Lou Y. 2011.** Genome-wide
930 transcriptional changes and defence-related chemical profiling of rice in
931 response to infestation by the rice striped stem borer *Chilo suppressalis*.
932 *Physiologia Plantarum* **143**: 21-40.

933 **Zipfel C. 2014.** Plant pattern-recognition receptors. *Trends in Immunology* **35**:
934 345-351.

935

936

Figure Legends

Fig. 1. Subcellular localization of OsLRR-RLK1.

Agrobacterium-mediated transient expression in *Nicotiana benthamiana* leaves of AtPIP2A-mCherry and OsLRR-RLK1-enhanced yellow fluorescent protein (EYFP). The first column shows mCherry fluorescence, and the second column shows the EYFP fluorescence. Overlaid image indicates co-localization of AtPIP2A-mCherry and OsLRR-RLK1-EYFP. White square in the overlaid image is shown as a detailed picture which is magnified in the fourth column. Yellow line in the detailed picture indicates the region of interest (ROI) that corresponds to the intensity profile in the last column. Intensity profile indicates the gray value of pixels across the ROI in the mCherry and EYFP channels. Leaf epidermal cells were imaged by confocal microscopy 72 h after infiltration with a suspension of each *Agrobacterium tumefaciens* strain at an OD₆₀₀ = 0.7. Scale bars: 20 µm.

Fig. 2. Transcriptional regulation of *OsLRR-RLK1*.

Mean transcript levels (+SE, $n = 5$) of *OsLRR-RLK1* in rice stems that were infested by rice striped stem borer (SSB, a), mechanically wounded (W, b), treated by *Spodoptera frugiperda* oral secretions (OS) after wounding (W + *S. frugiperda* OS, c), or jasmonic acid (JA, d). Con, control plants; Buf, buffer. Transcript levels were analyzed by quantitative real-time PCR. Asterisks represent significant differences between treatments and controls at the indicated times (Two-way analysis of variance [ANOVA], followed by pairwise comparisons of Least Squares Means [LSMeans], P values were corrected by False Discovery Rate [FDR] method; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Fig. 3. *OsLRR-RLK1* acts upstream of MPK cascades.

(a) MPK activation in *ir-lrr* lines and wild-type (WT) plants, which were infested by a third-instar striped stem borer (SSB) larva. Infested stems from five replicate plants were harvested at indicated times. Immunoblotting was performed using either anti-pTEpY antibody (upper panel) to detect phosphorylated MPKs, or actin antibody (lower panel) as a loading control which was detected on a replicate blot. For

quantification of immunodetection signals, see Fig. S10. This experiment was repeated three times, and the effect of *OsLRR-RLK1* was consistently observed (Fig. S10). (b to d) Mean transcript levels (+SE, $n = 5$) of *OsMPK3* (b), *OsMEK4* (c) and *OsMPK6* (d) in *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva. (e, f) Mean transcript levels (+SE, $n = 5$) of *OsLRR-RLK1* in *as-mpk3* (e), *as-mpk6* (f) and WT plants that were individually infested by a third-instar SSB larva. Asterisks represent significant differences between *ir-lrr* lines and WT plants at indicated times (Two-way analysis of variance [ANOVA], followed by pairwise comparisons of Least Squares Means [LSMeans], P values were corrected by False Discovery Rate [FDR] method; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Fig. 4. *OsLRR-RLK1* regulates defense-related WRKY transcription factors.

Mean transcript levels (+SE, $n = 5$) of *OsWRKY70* (a), *OsWRKY53* (b), *OsWRKY45* (c), *OsWRKY24* (d), *OsWRKY30* (e) and *OsWRKY33* (f) in *ir-lrr* lines and wild-type (WT) plants that were individually infested by a third-instar striped stem borer larva. Asterisks represent significant differences between *ir-lrr* lines and WT plants at indicated times (Two-way analysis of variance [ANOVA], followed by pairwise comparisons of Least Squares Means [LSMeans], P values were corrected by False Discovery Rate [FDR] method; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Fig. 5. *OsLRR-RLK1* mediates herbivore-induced jasmonic acid (JA), salicylic acid (SA) and ethylene (ET) biosynthesis.

(a, b) Mean levels (+SE, $n = 5$) of JA (a) and JA-Ile (b) in *ir-lrr* lines and wild-type (WT) plants that were individually infested by a third-instar striped stem borer (SSB) larva. (c, d) Mean transcript levels (+SE, $n = 5$) of *OsHI-LOX* (c) and *OsAOS1* (d) in *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva. (e) Mean levels (+SE, $n = 5$) of ET in *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva. (f) Mean transcript levels (+SE, $n = 5$) of *OsACS2* in *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva. (g) Mean levels (+SE, $n = 5$) of SA in *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva. (h) Mean transcript levels (+SE, $n = 5$) of SA biosynthesis-related gene *OsICS1* in *ir-lrr* lines and WT plants that were

individually infested by a third-instar SSB larva. FW, fresh weight. Asterisks represent significant differences between *ir-lrr* lines and WT plants at indicated times (Two-way analysis of variance [ANOVA], followed by pairwise comparisons of Least Squares Means [LSMeans], *P* values were corrected by False Discovery Rate [FDR] method; *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001).

Fig. 6. *OsLRR-RLK1* is not regulated by defense hormone signaling cascades.

Mean transcript levels (+SE, *n* = 5) of *OsLRR-RLK1* in *as-lox* (a), *as-aos1* (b), *as-ics1* (c), *as-acs2* (d) lines and wild-type (WT) plants that were individually infested by a third-instar striped stem borer larva.

Fig. 7. Silencing of *OsLRR-RLK1* attenuates trypsin protease inhibitor (TrypPI) activity and rice resistance to the striped stem borer (SSB).

(a) Mean TrypPI activity (+SE, *n* = 5) in *ir-lrr* lines and wild-type (WT) plants that were individually infested by a third-instar SSB larva for 3 days. (b) Mean larval weight (+SE, *n* = 30) of SSB feeding on *ir-lrr* lines or WT plants for 12 days. Letters indicate significant differences between lines (one way-analysis of variance [ANOVA], followed by multiple comparisons of Least Squares Means [LSMeans], which were corrected using False Discovery Rate [FDR] method, *P* < 0.05). (c) Mean activity (+SE, *n* = 5) of TrypPIs in *ir-1* line and WT plants which were individually treated with 100 µg of methyl jasmonate (MeJA) in 20 µl of lanolin paste (+MeJA) followed by a SSB larva feeding for 3 days. (d) Mean larval weight (+SE, *n* = 30) of SSB larvae 12 d after feeding on *ir-1* and WT plants that were individually treated with 100 µg of MeJA in 20 µl of lanolin paste (+MeJA). (e) Mean activity (+SE, *n* = 5) of TrypPIs in *ir-1* line and WT plants which were individually treated with 20 µl of pure lanolin paste (+Lanolin) followed by a SSB larva feeding for 3 days. (f) Mean larval weight (+SE, *n* = 30) of SSB larvae 12 d after feeding on *ir-1* and WT plants that were individually treated with 20 µl of pure lanolin paste (+Lanolin). Asterisks represent significant differences between *ir-1* and WT plants (Student's *t* tests, **, *P* < 0.01). (g) Damaged phenotypes of *ir-lrr* lines and WT plants that were individually infested by a third-instar SSB larva for 7 days (*n* = 20).

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Experimental setup used to infest rice plants with striped stem borer (SSB) larvae.

Fig. S2 Transformation vector used in this study.

Fig. S3 Nucleotide sequence and the deduced amino acid sequence of *OsLRR-RLK1*.

Fig. S4 Phylogenetic analysis of defense-related leucine rich repeat receptor-like kinases from Arabidopsis, tobacco and rice.

Fig. S5 Protein alignment of *OsLRR-RLK1* with homologous proteins in Arabidopsis.

Fig. S6 Salicylic acid (SA) treatment does not induce the expression of *OsLRR-RLK1*.

Fig. S7 DNA gel-blot analysis of *ir-lrr* and wild-type (WT) plants.

Fig. S8 Reduction of *OsLRR-RLK1* does not co-silence the transcript levels of its highly similar genes.

Fig. S9 Growth phenotypes of *ir-lrr* and wild-type (WT) plants.

Fig. S10 Relative activation of *OsMPK3* and *OsMPK6* in *ir-lrr* and wild-type (WT) plants.

Fig. S11 *OsLRR-RLK1* does not regulate wound-elicited *OsMPK3* and *OsMPK6* activation and the levels of JA and SA.

Table S1 Primers and probes used for QRT-PCR of target genes.